

Let's Take It From the Top

Ramon Diego, Haley Gipson, Huaidian Hou, and Anna Huang
Mentors: Dr. Alejandro Bravo-Doddoli and Kausik Das

University of Michigan Laboratory of Geometry



ABSTRACT

- The study of rigid body motion is equivalent to the study of dynamics in the group of rotations \mathbb{R}^3 denoted by $SO(3)$.
- General dynamics in $SO(3)$ are **chaotic**. There exist only three examples where the dynamics are **integrable** (non-chaotic): Lagrange, Euler and Kovalevskaya tops.
- These tops have many potential applications in the study of planetary motion, but they are difficult to visualize.

OBJECTIVE:
Simulate Lagrange, Euler, and Kovalevskaya Tops in MATLAB

THEORETICAL BACKGROUND

RIGID BODY MOTION: Tops (and also the Earth) can undergo three motions




Figure 1: Rotation, precession, and nutation diagram

LIE GROUPS


- Lie groups model continuous symmetries, e.g. $SO(3)$ is the group of rotations in \mathbb{R}^3 .
- The tangent space $T^*SO(3)$ = space of velocities where the Poisson bracket is defined.
- Poisson bracket $\{F, H\}$ is a tool we can use to find the equations of motion of our tops where F is a smooth function and H is the Hamiltonian.

HAMILTON'S EQUATIONS OF MOTION [Hamilton]

$$\dot{x}_i = \frac{\partial H}{\partial p_i}, \quad \text{and} \quad \dot{p}_i = -\frac{\partial H}{\partial x_i}$$

where $H(x, p)$ given $x = (x_1, \dots, x_n)$ is position and $p = (p_1, \dots, p_n)$ is momentum

EULER ANGLES



Definition. F is a **constant of motion** if its Poisson bracket commutes with H . A given H in $T^*SO(3)$ will be **integrable** if it has 3 linearly independent constants of motion that the Poisson bracket commutes between.

MOMENTS OF INERTIA

- Moment of inertia I = "rotational mass."
- There exists a natural orthonormal set of principal axes for a rigid body I_1, I_2 and I_3 that simplify equations of motion.

TOP ANIMATIONS




Figure 2: Euler Top Motion Snapshots




Figure 3: Lagrange Top Animation Interface Snapshot




Figure 4: Kovalevskaya Top Animation Interface Snapshot

| Type | Moments of Inertia | Constants of Motion | Hamiltonian H |
|--------------|--------------------|----------------------------|--|
| Lagrange | $I_1 = I_2 < I_3$ | p_ϕ, p_ψ and H | $\frac{p_\phi^2}{2I_1} + \frac{(p_\phi - p_\psi \cos(\theta))^2}{2I_1} + \frac{p_\psi^2}{2I_3} + gml \cos(\theta)$ |
| Euler | $I_1 < I_2 < I_3$ | p_1, p_2, p_3 | $\frac{I_1 \omega_1^2}{2} + \frac{I_2 \omega_2^2}{2} + \frac{I_3 \omega_3^2}{2}$ |
| Kovalevskaya | $I_1 = I_2 = 2I_3$ | p_θ, p_ψ and H | $\frac{p_\theta^2}{2I} + \frac{(p_\theta - p_\psi \cos(\theta))^2}{2I \sin^2(\theta)} + \frac{p_\psi^2}{4I} + gma \sin(\theta) \sin(\psi)$ |

Table 1: Tops' Properties Table

COMPUTATIONAL METHODS

- We derived differential equations for the Euler angle rates from our Hamiltonians

Example: Differential Equations for Lagrange Top

$$\frac{d\phi}{dt} = \frac{b - a \cos(\theta)}{\sin^2(\theta)}, \quad \frac{d\psi}{dt} = a \frac{I_1}{I_2} \frac{(b - a \cos(\theta)) \cos(\theta)}{\sin^2(\theta)},$$

$$\frac{d^2\theta}{dt^2} = \frac{(a^2 + b^2) \cos(\theta)}{\sin^2(\theta)} - ab \frac{3 + \cos(2\theta)}{2 \sin^3(\theta)} + \frac{\beta}{2} \sin(\theta)$$

- To find the equations of motion, we used ordinary differential equation solvers and other numerical integration methods.
- Our packages support custom initial conditions including angular velocity, moments of inertia, and initial heading. It also supports configuring multiple tops within one file.

RESULTS

EULER TOP VIDEO

- Gimbal locking causes momentary spike in the Euler angle rates.



LAGRANGE TOP VIDEO

- Nutation causes effective potential U_{eff} to oscillate.

KOVALEVSKAYA TOP VIDEO

- Back-and-forth motion: a combination of rotation and precession.



APPLICATIONS

- Integrable tops have many potential applications to planetary motion because planets are chaotic systems, making them difficult to study.
- Earth is similar to a Lagrange top, but its motion is closer to that of a **fast top**, i.e. it rotates much faster than it nutates or precesses.
- For reference, the Earth rotates roughly every 24 hours, nutates every 18.6 years, and precesses every 26,000 years!

FUTURE RESEARCH:
Study Connections Between Motion of Earth and Fast Tops

REFERENCES

- [1] Aleksei Viktorovich Bolsinov, PH Richter, and Anatoly Timofeevich Fomenko. The method of loop molecules and the topology of the kovalevskaya top. Sbornik: Mathe- matics, 191(2):151, 2000.
- [2] Illinois Geometry Lab. IGL Poster Template. *University of Illinois at Urbana-Champaign Department of Mathematics*, 2017.
- [3] John Robert Taylor and John R Taylor. Classical mechanics, volume 1. Springer, 2005.
- [4] John Stillwell. Naive lie theory. Springer Science & Business Media, 2008.
- [5] Lev Davidovich Landau and Evgeny Mikhailovich Lifshitz. Mechanics, volume 1. CUP Archive, 1960.

Thank you Dr. Alejandro Bravo-Doddoli, Kausik Das, and Dr. Nir Gadish for supporting our research project.